



On cloud modelling and the mass accommodation coefficient of water

A. Laaksonen, T. Vesala, M. Kulmala, P. M. Winkler, P. E. Wagner

► To cite this version:

A. Laaksonen, T. Vesala, M. Kulmala, P. M. Winkler, P. E. Wagner. On cloud modelling and the mass accommodation coefficient of water. *Atmospheric Chemistry and Physics Discussions*, 2004, 4 (6), pp.7281-7290. hal-00301497

HAL Id: hal-00301497

<https://hal.science/hal-00301497>

Submitted on 5 Nov 2004

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Cloud modelling and
accommodation
coefficient**

A. Laaksonen et al.

On cloud modelling and the mass accommodation coefficient of water

A. Laaksonen¹, T. Vesala², M. Kulmala², P. M. Winkler³, and P. E. Wagner³

¹Department of Applied Physics, University of Kuopio, P.O. Box 1627, FIN-70211 Kuopio, Finland

²Department of Physical Sciences, University of Helsinki, P.O. Box 64, FIN-00014 Helsinki, Finland

³Institut für Experimentalphysik, Universität Wien, Boltzmanngasse 5, A-1090 Wien, Austria

Received: 8 September 2004 – Accepted: 27 October 2004 – Published: 5 November 2004

Correspondence to: A. Laaksonen (ari.laaksonen@uku.fi)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2004

Abstract

The mass accommodation coefficient of water is a quantity for which different experimental techniques have yielded conflicting values in the range 0.04–1. From the viewpoint of cloud modelling, this is an unfortunate situation, since the value of the mass accommodation coefficient affects the model results, e.g. the number concentration of activated cloud droplets. In this paper we argue that a mass accommodation coefficient of unity should be used in cloud modelling, since this value has been obtained in experimental studies of water droplet growth rates, a quantity which is explicitly described in cloud models. In contrast, mass accommodation coefficient values below unity have been derived from experimental results which are analyzed with different theoretical expressions than those included in cloud models.

1. Introduction

The mass accommodation coefficient α of water vapor molecules on liquid water has been studied experimentally and theoretically for decades with conflicting results. Recent experiments on droplet growth rates (Winkler et al., 2004) indicate a mass accommodation coefficient of unity or near unity, while results from droplet train flow reactors (Li et al., 2001) have yielded values on the order of 0.1–0.3. Even lower values, on the order of 0.04–0.1, were obtained with a technique measuring droplet evaporation rates in an electrodynamic levitation chamber (Shaw and Lamb, 1999).

The mass accommodation coefficient is a quantity that affects among other things the results obtained with process models simulating cloud droplet growth. A mass accommodation coefficient below unity slows down droplet growth and causes the maximum supersaturation to increase. It has been shown that relatively large increases in cloud droplet number concentrations can result for small decreases in the value of α (Rudolf et al., 2001; Nenes et al., 2001). The values of mass accommodation coefficient applied in recent cloud model studies are between 0.04–1 (Kreidenweis et al.,

Cloud modelling and accommodation coefficient

A. Laaksonen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

2003), which undoubtedly causes differences in the model results. It is our purpose to point out that the droplet growth rate obtained in cloud model studies is only consistent with experimental results if a mass accommodation coefficient of unity is applied.

2. Discussion

- 5 The condensation theory used in cloud models is the so called transition regime condensation theory. The mass flux directed away from the droplet can be given in the form (Kulmala et al., 1993a,b)

$$I = \frac{-4\pi a(S - S_a)}{\frac{RT_\infty}{M_v \beta_M D(1+(S+S_a)p_{ve}(T_\infty)/2p)p_{ve}(T_\infty)} + \frac{S_a L^2 M_v}{R \beta_T K T_\infty^2}}, \quad (1)$$

- 10 where a is the droplet radius, S is the gas phase activity at ambient (far from the droplet) temperature T_∞ and S_a is the activity over the droplet surface, R is the gas constant, M_v is the molecular weight of the condensing vapour, D is the binary (between vapour and the inert gas) diffusion coefficient at T_∞ , p_{ve} is the saturation vapour pressure of the liquid, L is the latent heat of vaporization and K is the heat conductivity of the vapour-gas mixture at the ambient temperature and p is the ambient total gas pressure.
- 15 β_M is the transition regime correction for mass transfer (Fuchs and Sutugin, 1970)

$$\beta_M = \frac{1 + Kn}{1 + \left(\frac{4}{3\alpha_M} + 0.377\right)Kn + \frac{4}{3\alpha_M}Kn^2}, \quad (2)$$

where the Knudsen number Kn is the ratio of the mean effective free path of the vapour molecules, calculated from the vapour diffusivity, to the droplet radius. α_M is now the mass accommodation coefficient. Correspondingly, β_T is the transitional correction

factor for heat transfer

$$\beta_T = \frac{1 + Kn_T}{1 + \left(\frac{4}{3\alpha_T} + 0.377 \right) Kn_T + \frac{4}{3\alpha_T} Kn_T^2}, \quad (3)$$

where α_T is the thermal accommodation coefficient. The Knudsen number Kn_T for heat transfer is defined analogously to Kn by replacing the mean free path of vapour by a length scale for heat transfer, which is provided by the mean effective free path of the carrier gas molecules calculated from the heat conductivity of the inert gas (Wagner, 1982). Clearly, α_M refers to water vapour molecules, while α_T refers (mainly) to inert air molecules. Note, that the transition regime corrections used in the present study in conjunction with the proper definitions of the Knudsen numbers (Fuchs and Sutugin, 1970) have been found to provide good approximations for molecular mass ratios ranging from values $\ll 1$ (light vapours) up to values exceeding 10 (Qu and Davis, 2001). Similar expressions must also be applied in multicomponent condensation calculations, see e.g. the treatment of nitric acid condensation during cloud formation by Kulmala et al. (1993b).

In calculations of the condensational growth, the equations for mass and heat fluxes are coupled and knowledge of the droplet temperature is required in order to calculate the mass flux. The above expression for the mass flux takes approximately into account the correct droplet temperature due the latent heat release and under atmospheric conditions the formula very precisely gives the same results as the full coupled equations used by Winkler et al. (2004) (see also Vesala et al., 1997). Furthermore, Fladerer et al. (2002) have shown that the formula is applicable to all growth regimes and to initial conditions of high supersaturation, for which it was not expected to work properly.

Winkler et al. (2004) carried out experiments on the growth rates of water droplets observing growth kinetics in an expansion cloud chamber system. They studied liquid droplets nucleated on Ag particles and growing due to condensation of supersaturated water vapour using the experimental system presented in detail by Wagner

Cloud modelling and accommodation coefficient

A. Laaksonen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

et al. (2003). Vapour supersaturation is achieved by adiabatic expansion in a computer controlled thermostated expansion chamber resulting in well defined uniform thermodynamical conditions in the measuring volume. Growth of droplets is observed using the constant-angle Mie scattering (CAMS) detection method (Wagner, 1985) providing absolute, time-resolved and non-invasive simultaneous determination of droplet diameter and number density. Winkler et al. (2004) compared their results with theoretical calculations employing the transition regime condensation correction by Fuchs and Sutugin (1970) with different values of the mass accommodation coefficient. They found agreement only for mass accommodation coefficients in a certain range around unity.

The droplet train apparatus of Li et al. (2001) is based on a fast-moving monodisperse, spatially collimated train of droplets interacting with the gas-phase species (H_2^{17}O or D_2O) in trace quantities. The liquid water itself is in equilibrium with its vapor, and the uptake of the trace isotopic species (which is of course out-of equilibrium) does not significantly perturb the bulk phase or the surface of the liquid. Mass accommodation coefficients are obtained from the determination of uptake (condensation) of gas phase isotope in trace amounts combined with calculations using a theory which describes the transport of vapor molecules from the gas phase to the air-water interface, and transfer of the species across the interface, i.e. a different theory than that used by Winkler et al. (2004), and in the cloud models. Attempts have been made to reconcile the above results (Morita et al., 2004, Davidovits et al., submitted, 2004¹) but so far without success. Both the growth rate and droplet train experiments are carefully designed, and almost an order of magnitude difference in the mass accommodation coefficient seems quite high.

The lowest experimental values measured for α in recent years, between 0.04–0.1, were obtained by Shaw and Lamb (1999). They measured the evaporation rate of droplets suspended in an electrodynamic levitation system within a controlled envi-

¹Davidovits, P., Worsnop, D. R., Jayne, J. T., Kolb, C. E., Winkler, P., Vrtala, A., Wagner, P. E., Kulmala, M., Lehtinen, K. E. J., Vesala, T., and Mozurkewich, M.: Mass Accommodation coefficient of water vapor on liquid water, J. Geophys. Res., submitted, 2004.

Cloud modelling and accommodation coefficient

A. Laaksonen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

ronment. They used the homogeneous freezing nucleation rate as a way to measure droplet temperatures (Pruppacher and Klett, 1997). However, this technique might be prone to errors since the homogeneous nucleation theory is still a complex, unsolved problem (see e.g. Seinfeld and Pandis, 1998).

Figure 1 shows the experimental results of Winkler et al. (2004) for droplet growth, as well as theoretical growth curves calculated using the transition regime condensation theory, and mass accommodation coefficients ranging from 0.04 to 1. It is clear that only values near unity yield theoretical predictions consistent with the growth rates. Accordingly, the full coupled droplet growth equations accurately predict observations of droplet growth kinetics in an expansion cloud chamber system (Winkler et al., 2004), when the accommodation coefficients α_M and α_T are set to 1. Thus the usage of expressions (1–3) with α_M and $\alpha_T=1$ is a physically rigorous and consistent approach to estimate the condensational growth. We stress rigor, since many atmospheric scientists are apparently unaware of the recent developments in the condensational growth theories and apply fundamentally incorrect expressions deviating from that given above, and very recently discussed by Pines et al. (2004) and Vesala et al. (2004). We stress consistency, since the same theory used in the interpretations of the experiments must be fully applied in subsequent atmospheric models. Therefore, as long as the transition regime condensation theory is used in cloud models, the accommodation coefficients should be set to unity. Lower values can lead to too high cloud drop number concentrations, as indicated in Table 1.

3. Conclusions

At present, it is not clear whether the real value of α is near 1 or below. If the latter case proves to be correct, this must be associated with a deficiency of the currently used transition regime corrections for droplet growth rates. Whatever the case, the rigorous transition regime growth theory combined with a water vapour thermal and mass accommodation coefficients of unity yields excellent predictions of experimental

Cloud modelling and accommodation coefficient

A. Laaksonen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

droplet growth rates. For this reason, the use of mass accommodation coefficient values lower than 1 in cloud models together with the rigorous droplet growth theory is inconsistent and should be avoided.

Acknowledgements. This work was supported by the Academy of Finland (project 201052).

References

- Fladerer, A., Kulmala, M., and Strey, R.: Test of the applicability of Kulmala's analytical expression for the mass flux of growing droplets in highly supersaturated systems: growth of homogeneously nucleated water droplets, *J. Aerosol. Sci.*, 33, 391–399, 2002. [7284](#)
- Fuchs, N. A. and Sutugin, A. G.: Highly Dispersed Aerosols, Ann Arbor Science Publ., Ann Arbor, Michigan, 1970. [7283](#), [7284](#), [7285](#), [7290](#)
- Kreidenweis, S. M., Walcek, C. J., Feingold, G., Gong, W., Jacobson, M. Z., Kim, C.-H., Liu, X., Penner, J. E., Nenes, A., and Seinfeld, J. H.: Modification of aerosol mass and size distribution due to aqueous-phase SO₂ oxidation in clouds: Comparison of several models, *J. Geophys. Res.*, 108 (D7), 4213, doi:10.1029/2002JD002697, 2003. [7282](#)
- Kulmala, M. and Wagner, P. E.: Mass accommodation and uptake coefficients – a quantitative comparison, *J. Aerosol Sci.*, 32, 833–841, 2001.
- Kulmala, M., Vesala, T. and Wagner, P. E.: An analytical expression for the rate of binary condensational particle growth, *Proc. R. Soc. Lond. A*, 441, 589–605, 1993a. [7283](#)
- Kulmala, M., Laaksonen, A., Korhonen, P., Vesala, T., Ahonen, T., and Barrett, J. C.: The effect of atmospheric nitric acid vapor on cloud condensation nucleus activation, *J. Geophys. Res.*, 98 (D12), 22 949–22 958, 1993b. [7283](#), [7284](#)
- Li, J. L., Shi, O., Davidovits, P., Worsnop, D. R., Jayne, J. T., and Kolb, C. E.: Mass and Thermal Accommodation Coefficients of H₂O(g) on Liquid Water as a Function of Temperature, *J. Phys. Chem. A*, 105, 10 627–10 634, 2001. [7282](#), [7285](#)
- Morita, A., Sugiyama, M., Kameda, H., Koda, S., and Hanson, D. R.: Mass accommodation coefficient of water: molecular dynamics simulation and revised analysis of droplet train/flow reactor experiment, *J. Phys. Chem. B*, 108, 9111–9120, 2004. [7285](#)
- Nenes, A., Ghan, S., Abdul-Razzak, H., Chuang, P. Y., and Seinfeld, J. H.: Kinetic limitations on cloud droplet formation and impact on cloud albedo, *Tellus*, 53B, 133–149, 2001. [7282](#)

Cloud modelling and accommodation coefficient

A. Laaksonen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

- Pines, V., Zlatkowski, M., and Chait, A.: Postnucleation droplet growth in supersaturated gas with arbitrary vapor concentration, *J. Chem. Phys.*, 10 455–10 469, 2004. [7286](#)
- Pruppacher, H. R. and Klett, J. D.: *Microphysics of clouds and precipitation*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1997. [7286](#)
- 5 Rudolf, R., Vrtala, A., Kulmala, M., Vesala, T., Viisanen, Y., and Wagner, P. E.: Experimental study of sticking probabilities for condensation of nitric acid – water vapor mixtures, *J. Aerosol Sci.*, 32, 913–932, 2001. [7282](#)
- Qu, X. and Davis, A. J.: Droplet evaporation and condensation in the near-continuum regime, *J. Aerosol Sci.*, 32, 861–876, 2001. [7284](#)
- 10 Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics*, John Wiley and Sons, New York, 1998. [7286](#)
- Shaw, R. A. and Lamb, D.: Experimental determination of the thermal accommodation and condensation coefficients of water, *J. Chem. Phys.*, 111, 10 659–10 663, 1999. [7282](#), [7285](#)
- Vesala, T., Kulmala, M., Rudolf, R., Vrtala, A., and Wagner, P. E.: Models for condensational growth and evaporation of binary aerosol particles, *J. Aerosol Sci.*, 28, 565–598, 1997. [7284](#)
- 15 Vesala, T., Kulmala, M., and Wagner, P. E.: Comment on “Postnucleation droplet growth in supersaturated gas with arbitrary vapor concentrations”, (*J. Chem. Phys.*, 120, 10 455, 2004), *J. Chem. Phys.*, 121, 8163–8164, 2004. [7286](#)
- Wagner, P. E.: Aerosol growth by condensation, in *Aerosol Microphysics II.*, edited by: Marlow, W. H., Springer-Verlag, Düsseldorf, Germany, 129–178, 1982. [7284](#)
- 20 Wagner, P. E.: A constant-angle mie scattering method (cams) for investigation of particle formation processes, *J. Colloid Interface Sci.*, 105, 456–467 1985. [7285](#)
- Wagner, P. E., Kaller, D., Vrtala, A., Lauri, A., Kulmala, M., and Laaksonen, A.: Nucleation probability in binary heterogeneous nucleation of water-n-propanol vapor mixtures on insoluble and soluble nanoparticles, *Phys. Rev. E*, 67, 021605, 2003. [7284](#)
- 25 Winkler, P. M., Vrtala, A., Wagner, P. E., Kulmala, M., Lehtinen, K. E. J., and Vesala, T.: Mass and thermal accommodation during gas-liquid condensation of water, *Phys. Rev. Lett.*, 93, 075701, 2004.
- [7282](#), [7284](#), [7285](#), [7286](#)

Cloud modelling and accommodation coefficient

A. Laaksonen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Cloud modelling and accommodation coefficient

A. Laaksonen et al.

Table 1. The effect of accommodation coefficient on cloud drop number concentration (CDNC) calculated using a cloud parcel model. The aerosol size distribution is lognormal, with a mean diameter of 50 nm, standard deviation of 1.7, and number concentration of 3000 cm⁻³. The updraft velocity is 1.0 m/s and temperature 273 K.

α	CDNC (cm ⁻³)
0.04	1595
0.1	1221
1	873

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**Cloud modelling and
accommodation
coefficient**

A. Laaksonen et al.




Fig. 1. Experimental results of Winkler et al. (2004) for droplet growth. Theoretical growth curves calculated by means of the transitional drop growth theory (Fuchs and Sutugin, 1970) for mass accommodation coefficients between 0.04 and 1.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)